

# DOES TREATMENT EXPERIENCE MATTER?

Learn more about what  
once-monthly SIMPONI may  
be able to do for you

SIMPONI is a prescription medicine for adults with moderate to severe rheumatoid arthritis (RA), with the medicine methotrexate (MTX), active psoriatic arthritis (PsA), alone or with the medicine MTX, or active ankylosing spondylitis (AS).

---

SIMPONI 50 mg is given as an injection under the skin (subcutaneous injection) once a month.

MTX is used as directed.

Once you and your doctor are comfortable with the self-injection process, you will inject SIMPONI under the skin.

## **SELECTED IMPORTANT SAFETY INFORMATION**

SIMPONI<sup>®</sup> (golimumab) can lower your ability to fight infections. Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma, other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages [12-14](#) and the [Medication Guide](#), and talk with your doctor.

## WHAT CAN YOU EXPECT FROM YOUR SIMPONI EXPERIENCE?

When you live with the symptoms of RA, PsA, or AS, it's important to find the medication that is right for you. SIMPONI may help relieve the symptoms of moderate to severe RA, with the medicine MTX, active PsA, alone or with the medicine MTX, or active AS.

### HOW CAN SIMPONI HELP?

In clinical trials, one injection of SIMPONI monthly has been proven to help relieve:

- joint pain, stiffness, and swelling of moderate to severe RA
- joint pain, stiffness, swelling, and skin symptoms of active PsA. SIMPONI is not approved to treat plaque psoriasis
- pain and stiffness of active AS

Results may not be the same for everyone.



### SELECTED IMPORTANT SAFETY INFORMATION

SIMPONI® (golimumab) can lower your ability to fight infections. Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma,



"SIMPONI, taken with methotrexate, has helped me be active again by reducing the pain and swelling in my joints from RA. I can climb up the stairs in my house, walk my dog, and put more work into my garden."\*

MTX is taken as directed. Your results may vary. Talk with your doctor about your response to SIMPONI.

\*Based on patient stories. Not an actual patient quote or image.

### SIMPONI ONCE-MONTHLY EXPERIENCE

SIMPONI is the first once-monthly self-injectable biologic treatment with just one injection each month for adults with:

- moderate to severe RA, with the medicine MTX
- active PsA, alone or with the medicine MTX
- active AS

For RA, PsA, and AS: SIMPONI 50 mg is given as an injection under the skin (subcutaneous injection) once a month.

MTX is used as directed. Once you and your doctor are comfortable with the self-injection process, you will inject SIMPONI under the skin just once a month.

Your doctor should test you for tuberculosis and hepatitis B before starting SIMPONI.

other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages 12-14 and the Medication Guide, and talk with your doctor.



## ONCE-MONTHLY DOSING DESIGNED WITH PATIENTS IN MIND

Minimum number of doses first year.

**12**  
DOSES




**SIMPONI** autoinjector + MTX

SIMPONI is indicated for the treatment of adult patients with moderately to severely active RA, in combination with the medicine MTX.

MTX=methotrexate.


**26**  
DOSES



**Humira**® (adalimumab) pen

Humira® is indicated for reducing signs and symptoms, inducing major clinical response, inhibiting the progression of structural damage, and improving physical function in adult patients with moderately to severely active RA. Humira® can be used alone or in combination with MTX or other non-biologic disease-modifying anti-rheumatic drugs (DMARDs).

**52**  
DOSES



**Enbrel**® (etanercept) autoinjector

Enbrel® is indicated for reducing signs and symptoms, inducing major clinical response, inhibiting the progression of structural damage, and improving physical function in patients with moderately to severely active RA. Enbrel® can be initiated in combination with MTX or used alone.

TNF-alpha inhibitors used under the skin (subcutaneous injection) in a clinical study were Humira® and Enbrel®.

\*Trademarks are the property of their respective owners.  
This presentation is not intended to compare the safety, effectiveness, or uses of these treatments. This information, along with other factors, can be considered with your doctor when discussing treatment options. Please refer to each product's Prescribing Information and Medication Guide for recommended dosing and administration.

While dosing may be important, there are other factors to consider. So talk to your doctor about treatment options and what might be right for you.

### SELECTED IMPORTANT SAFETY INFORMATION

**SIMPONI® (golimumab) can lower your ability to fight infections.** Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma,

## MEET THE SMARTJECT® AUTOINJECTOR DESIGNED FOR AN EASY-TO-USE EXPERIENCE



- HIDDEN NEEDLE**  
Thin needle remains hidden throughout the injection process so you never have to see it
- GREEN SAFETY SLEEVE**  
Helps ensure that you'll be able to inject your medicine only when the autoinjector is firmly pressed against your skin
- ACTIVATION BUTTON**  
A simple squeeze of the side activation button injects SIMPONI under the skin
- EASY-TO-GRIP SHAPE**  
Oval-shaped body fits comfortably in your hand

*For those who prefer a prefilled syringe, SIMPONI gives you that choice too. As with the SmartJect® autoinjector, the first time you use it should be in your doctor's office with the help of your doctor or nurse.*



*Both the SmartJect® autoinjector and prefilled syringe have been awarded the Ease-of-Use® Commendation by the Arthritis Foundation.*

If your doctor decides that you or your caregiver may be able to give your injections at home, you should receive training on the right way to prepare and inject SIMPONI. Do not try to inject SIMPONI® yourself until you have been shown the right way to give the injections by your doctor or nurse.

other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages 12-14 and the Medication Guide, and talk with your doctor.



## WHAT TO EXPECT FROM YOUR INJECTION EXPERIENCE WITH SIMPONI

IN CLINICAL STUDIES FOR RA, PsA, AND AS:

**6%**

**of patients treated with SIMPONI experienced injection site reactions**

6% of patients experience injection site reactions (vs 2% of patients taking placebo) such as redness, swelling, itching, pain, bruising, or tingling.

**<1%**

**of patients experienced any injection site pain**

Less than 1% of patients treated with either SIMPONI or placebo reported experiencing any injection site pain.

### SELECTED IMPORTANT SAFETY INFORMATION

SIMPONI® (golimumab) can lower your ability to fight infections. Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma,

## WHEN WILL YOU EXPERIENCE RELIEF FROM YOUR RA SYMPTOMS?

Patients using SIMPONI may experience symptom relief after 3 months. Studies show some patients saw that SIMPONI helped relieve the symptoms of moderate to severe RA, with the medicine MTX as directed, in just 1 month.

LET'S GET STARTED

**VISIT [SIMPONI.COM](http://SIMPONI.COM) TO LEARN MORE ABOUT WHAT YOU CAN EXPECT FROM YOUR SIMPONI EXPERIENCE.**

other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages 12-14 and the Medication Guide, and talk with your doctor.



## GETTING STARTED WITH J&J withMe

### Patient Support for SIMPONI

Once you and your doctor have decided that SIMPONI is right for you, J&J withMe will help you find the resources you may need to get started and stay on track.

## J&J withMe

### Paying for SIMPONI

We can identify cost support options that may help with managing your out-of-pocket costs—whether you have commercial or private health insurance, government coverage such as Medicare or Medicaid, or have no insurance coverage.

### Getting Started

J&J withMe can review your health plan benefits and insurance coverage for SIMPONI, and offer treatment education resources.

### Staying on Track

We understand how important it is for you to take the SIMPONI your doctor prescribed. J&J withMe provides ongoing support to help you stay on track with your SIMPONI treatment.

## SELECTED IMPORTANT SAFETY INFORMATION

SIMPONI® (golimumab) can lower your ability to fight infections. Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma,

other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages 12-14 and the Medication Guide, and talk with your doctor.



**Call a Case Manager  
at 877-227-3728**

Monday – Friday, 8:00 AM to 8:00 PM ET.

**Set up a personal J&J withMe Account at  
[Account.JNJwithMe.com](https://www.JNJwithMe.com)** where you can:

check your insurance coverage for SIMPONI; if eligible, enroll in the J&J withMe Savings Program; and sign up for treatment support.

Visit [JNJwithMe.com/Simponi](https://www.JNJwithMe.com/Simponi)



## GET SUPPORT FOR YOUR MEDICINE FROM JOHNSON & JOHNSON

### Paying for SIMPONI

At J&J, we don't want cost to get in the way of treatment you need.

We can help you explore options to lower your out-of-pocket cost for SIMPONI.

No matter what type of insurance you have—or even if you don't have insurance—J&J withMe can help explain your medicine insurance coverage and benefits, and help find programs that may help you pay for SIMPONI.

If you have commercial or private health insurance and need help paying for SIMPONI, the J&J withMe Savings Program may be able to help. For more information, visit [Simponi.JNJwithMeSavings.com](https://www.Simponi.JNJwithMeSavings.com) or call **J&J withMe at 877-227-3728**

If you don't have commercial or private health insurance, you may find help from the programs and resources found on [JNJwithMe.com](https://www.JNJwithMe.com)

### SELECTED IMPORTANT SAFETY INFORMATION

SIMPONI® (golimumab) can lower your ability to fight infections. Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma,

## J&J withMe SAVINGS PROGRAM

\*Eligible patients using commercial or private insurance can save on out-of-pocket medicine costs for SIMPONI.

### ELIGIBLE PATIENTS

**PAY \$5**

### FOR EACH INJECTION

Maximum program benefit per calendar year shall apply. Not valid for patients using Medicare, Medicaid, or other government-funded programs to pay for their medications. Terms expire at the end of each calendar year. Offer subject to change or end without notice. Restrictions, including monthly maximums, may apply. Program does not cover the cost to give you your injection. See program requirements at [Simponi.JNJwithMeSavings.com](https://www.Simponi.JNJwithMeSavings.com).

### Check your eligibility and enroll in the J&J withMe Savings Program

Create an online account and enroll at [Account.JNJwithMe.com](https://www.Account.JNJwithMe.com) or call a Case Manager at **877-227-3726**, Monday–Friday, 8:00 AM to 8:00 PM ET



"As a patient living with PsA, one of the things that makes SIMPONI more affordable for me is the J&J withMe Savings Program."\*

\*Based on patient stories. Not an actual patient quote or image.

other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages 12–14 and the Medication Guide, and talk with your doctor.

SUPPORT



## IMPORTANT SAFETY INFORMATION

### SERIOUS INFECTIONS

**SIMPONI® (golimumab) is a prescription medicine. SIMPONI® can lower your ability to fight infections. There are reports of serious infections caused by bacteria, fungi, or viruses that have spread throughout the body, including tuberculosis (TB) and histoplasmosis. Some of these infections have been fatal. Your doctor will test you for TB before starting SIMPONI® and will monitor you for signs of TB during treatment. Tell your doctor if you have been in close contact with people with TB. Tell your doctor if you have been in a region (such as the Ohio and Mississippi River Valleys and the Southwest) where certain fungal infections like histoplasmosis or coccidioidomycosis are common.**

You should not start SIMPONI® if you have any kind of infection. Tell your doctor if you are prone to or have a history of infections or have diabetes, HIV, or a weak immune system. You should also tell your doctor if you are currently being treated for an infection or if you have or develop any signs of an infection such as:

- fever, sweat, or chills
- muscle aches
- cough
- shortness of breath
- blood in phlegm
- weight loss
- warm, red, or painful skin or sores on your body
- diarrhea or stomach pain
- burning when you urinate or urinate more than normal
- feel very tired

Your doctor will examine you for TB and perform a test to see if you have TB. If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with SIMPONI® and during treatment with SIMPONI®. Even if your TB test is negative, your doctor should carefully monitor you for TB infections while you are taking SIMPONI®. People who had a negative TB skin test before receiving SIMPONI® have developed active TB. Tell your doctor if you have any of the following symptoms while taking or after taking SIMPONI®:

- cough that does not go away
- low grade fever
- weight loss
- loss of body fat and muscle (wasting)

### CANCER

**Unusual cancers have been reported in children and teenage patients taking TNF-blocker medicines.** For children and adults taking TNF blockers, including SIMPONI®, the chances for getting lymphoma or other cancers may increase. Hepatosplenic T-cell lymphoma, a rare and fatal lymphoma, has occurred mostly in teenage or young adult males with Crohn's disease or ulcerative colitis who were taking other TNF blockers with azathioprine or 6-mercaptopurine. You should tell your doctor if you have had or develop lymphoma or other cancers.

Some people treated with SIMPONI® have developed certain kinds of skin cancer. If any changes in the appearance of your skin or growths on your skin occur during or after your treatment with SIMPONI®, tell your doctor.

### USE WITH OTHER DRUGS

Tell your doctor about all the medications you take including ORENCIA® (abatacept), KINERET® (anakinra), ACTEMRA® (tocilizumab),

RITUXAN® (rituximab), or another TNF blocker, or if you are scheduled to or recently received a vaccine. People taking SIMPONI® should not receive live vaccines or treatment with a weakened bacteria (such as BCG for bladder cancer).

### HEPATITIS B INFECTION

If you are a carrier of the hepatitis B virus (a virus that affects the liver), the virus can become active while you use SIMPONI®. Some of these cases have been fatal. Your doctor should do blood tests before and while you are using SIMPONI®. Tell your doctor if you know or think you may be a carrier of hepatitis B virus or if you experience signs of hepatitis B infection, such as:

- feel very tired
- dark urine
- skin or eyes look yellow
- little or no appetite
- vomiting
- muscle aches
- clay-colored bowel movements
- fevers
- chills
- stomach discomfort
- skin rash

### HEART FAILURE

**Heart failure, including new heart failure or worsening of heart failure that you already have, can occur or get worse in people who use TNF blockers, including SIMPONI®.** If you develop new or worsening heart failure with SIMPONI®, you may need treatment in a hospital, and it may result in death. If you have heart failure before starting SIMPONI®, your condition should be watched closely during treatment with SIMPONI®. Call your doctor right away if you get new or worsening symptoms of heart failure like shortness of breath, swelling of your lower legs or feet, or sudden weight gain.

### NERVOUS SYSTEM PROBLEMS

Rarely, people using TNF blockers, including SIMPONI®, can have nervous system problems such as multiple sclerosis or Guillain-Barré syndrome. Tell your doctor right away if you have symptoms like vision changes, weakness in your arms or legs, or numbness or tingling in any part of your body.

### IMMUNE SYSTEM PROBLEMS

Rarely, people using TNF blockers have developed lupus-like symptoms. Tell your doctor if you have any symptoms such as a rash on your cheeks or other parts of the body, sensitivity to the sun, new joint or muscle pain, becoming very tired, chest pain or shortness of breath, or swelling of the feet, ankles, and/or legs.

### LIVER PROBLEMS

Serious liver problems can happen in people using TNF blockers, including SIMPONI®. These problems can lead to liver failure and death. Call your doctor immediately if you develop symptoms such as feeling very tired, skin or eyes look yellow, poor appetite or vomiting, or pain on the right side of your stomach (abdomen).

### BLOOD PROBLEMS

Low blood counts have been seen with SIMPONI®. If this occurs, your body may not make enough blood cells to help fight infections or help stop bleeding. Your doctor will check your blood counts before and during treatment. Tell your doctor if you have signs such as fever, bruising or bleeding easily, or looking pale.

Continued on next page.



## IMPORTANT SAFETY INFORMATION

Continued from previous page.

### ALLERGIC REACTIONS

Allergic reactions can happen in people who use TNF-blocker medicines, including SIMPONI®. Some reactions may be serious and can be life-threatening. Some of these reactions can happen after receiving your first dose of SIMPONI®. Stop using SIMPONI® and call your doctor right away if you have any symptoms of an allergic reaction such as hives, swollen face, breathing trouble, or chest pain.

### OTHER CONSIDERATIONS TO TELL YOUR DOCTOR

Tell your doctor if you are allergic to rubber or latex. The needle cover on the prefilled syringe and SmartJect autoinjector contains dry natural rubber.

Tell your doctor if you are pregnant, planning to become pregnant, breastfeeding, plan to breastfeed, or have a baby and were using SIMPONI® during pregnancy. Tell your baby's doctor before your baby receives any vaccine because of an increased risk of infection for up to 6 months after birth.

**The most common side effects of SIMPONI include:** upper respiratory infection (runny nose, sore throat, and hoarseness or laryngitis), reaction at site of injection (redness, swelling, itching, pain, bruising, or tingling), and viral infections such as flu and oral cold sores.

### PSORIASIS

New or worsening psoriasis symptoms may occur. Tell your doctor if you develop red scaly patches or raised bumps that are filled with pus. Your doctor may decide to stop your treatment with SIMPONI®.

These are not all of the possible side effects of SIMPONI®. Tell your doctor about any side effect that bothers you or does not go away. Call your doctor for medical advice about side effects.

**Please read the full Prescribing Information and Medication Guide for SIMPONI® and discuss any questions you have with your doctor.**

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.**

cp-51591v5

**GET STARTED WITH THE  
SIMPONI TREATMENT  
EXPERIENCE**

**JUST ONE INJECTION,  
ONCE A MONTH**

For more information  
visit [Simponi.com](http://Simponi.com) or call  
**877-227-3728**

For adults with moderate to severe RA, with the medicine MTX, active PsA, alone or with the medicine MTX, or active AS.

### SELECTED IMPORTANT SAFETY INFORMATION

SIMPONI® (golimumab) can lower your ability to fight infections. Serious and sometimes fatal events may occur. There have been reports of serious infections including tuberculosis (TB) and infections caused by bacteria, fungi, or viruses that have spread throughout the body. Other possible serious side effects may include lymphoma, skin cancer, a rare and fatal cancer called hepatosplenic T-cell lymphoma, other cancers, hepatitis B, heart failure, nervous system problems, immune system problems (lupus-like syndrome), liver problems, blood problems, or allergic reactions. To learn more about these and other risks, please read the Important Safety Information on pages 12-14 and the Medication Guide, and talk with your doctor.

© Johnson & Johnson and its affiliates 2026 06/26 cp-576630v1